arXiv:1209.0160v1 [math.CO] 2 Sep 2012

STRONGLY EVEN CYCLE DECOMPOSABLE GRAPHS

TONY HUYNH, ANDREW D. KING, SANG-IL OUM, AND MARYAM VERDIAN-RIZI

Abstract. We introduce the notion of an Eulerian graph being *strongly even cycle decomposable*, and note that a few basic classes have this property. We then prove that several fundamental composition operations that preserve the property of being Eulerian also preserve the property of being strongly even cycle decomposable. Our composition operations imply that all simple Eulerian k-partite graphs are strongly even cycle decomposable, with the exception of K_5.

1. Introduction

A graph G is *even cycle decomposable (ECD)* if the edge set of G can be partitioned into cycles of even length. Even cycle decomposable graphs have been the subject of substantial attention. For a summary of relevant results we refer to the reader to the surveys of Jackson [3] or Fleischner [1]. In addition, Huynh, Oum and Verdian-Rizi [2] recently proposed a conjecture that unifies many known results on ECD graphs.

In this paper we instead focus on a stronger decomposition property. Namely, we define a graph G to be *strongly even cycle decomposable (strongly ECD)* if every subdivision of G with an even number of edges is even cycle decomposable. Note that strongly even cycle decomposable graphs are necessarily Eulerian and 2-connected (after removing isolated vertices).

One motivation for introducing strongly even cycle decomposable graphs is that inductive arguments tend to work more smoothly for strongly ECD graphs as opposed to ECD graphs. For this reason, sometimes the easiest way to prove that a graph is even cycle decomposable is to prove that it is strongly even cycle decomposable. In addition, there are indeed interesting classes of graphs that are strongly even cycle decomposable. For example, Seymour [5] proved that planar graphs that satisfy the obvious necessary conditions are strongly even cycle decomposable.

Theorem 1.1 (Seymour [5])**.** *Every loopless 2-connected Eulerian planar graph is strongly ECD.*

Theorem 1.1 was generalized by Zhang [6] to graphs having no K_5-minor.

Theorem 1.2 (Zhang [6])**.** *Every loopless 2-connected Eulerian K_5-minor-free graph is strongly ECD.*

Date: September 2, 2012.

Key words and phrases. cycle, even cycle decomposition, Eulerian.

T. H., S. O., and M. V.-R. are supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012-0004119). S. O. is also supported by TJ Park Junior Faculty Fellowship. A. D. K. is supported by an EBCO/Ebbich Postdoctoral Scholarship and the NSERC Discovery Grants of Pavol Hell and Bojan Mohar.

We note that K_5 is an example where the obvious necessary conditions are not sufficient. To see this, observe that K_5 has 10 edges but every even length cycle of K_5 is a 4-cycle. Thus, K_5 is not ECD (and hence not strongly ECD). Markström [4] recently gave a construction for an infinite class of 4-regular 2-connected graphs that are not ECD; the construction is based on a gadget that embeds K_5 in an edge.

In this paper we prove that Eulerian complete bipartite graphs are strongly ECD, and that several fundamental composition operations that preserve the property of being Eulerian also preserve the property of being strongly ECD. In doing so we provide evidence that in some sense, K_5 seems to be the unique obstruction to the strongly ECD property. In particular, we obtain the following theorem as an easy corollary of our results.

Theorem 1.3. *All simple Eulerian complete k-partite graphs are strongly ECD, except for K_5.*

Using Theorem 1.1 or Theorem 1.2 to provide base classes for our composition operations, we also obtain numerous strongly ECD graphs that were previously not known to be strongly ECD.

The rest of the paper is organized as follows. In Section 2, we define *signed graphs* and recast our problem in their language. In Section 3 we prove that Eulerian complete bipartite graphs are strongly ECD. In Sections 4, 5, 6, 7, and 8 we introduce our composition operations and prove that they preserve the strongly ECD property. Finally, in Section 9 we derive Theorem 1.3 as an easy corollary of our composition operations.

2. Signed graphs

A *signed graph* is a pair (G, Σ) consisting of a graph G together with a *signature* $\Sigma \subseteq E(G)$. The edges in Σ are *odd* and the other edges are *even*. We abuse terminology and define a cycle (respectively, path) in a signed graph to be *even* if it contains an even number of *odd* edges. To avoid confusion, we will always use the term *even length cycle* (respectively, *even length path*) if we need to refer to a cycle (respectively, path) with an even number of edges. Odd length paths and cycles are defined accordingly. A signed graph (G, Σ) is *even cycle decomposable (ECD)*, if $E(G)$ can be partitioned into even cycles.

The relationship between even cycle decompositions of signed graphs and graphs is as follows. Let G be a graph and let H be a subdivision of G with $|E(H)|$ even. Consider the signed graph $\mathcal{G}_H := (G, \Sigma)$, where $e \in \Sigma$ if and only if the subdivided edge in H corresponding to e has odd length. Observe that H is ECD if and only if $\mathcal{G}_H$ is ECD. Thus, we have the following equivalent definition of strongly ECD graphs.

Definition 2.1. A graph G is strongly ECD if and only if for each signature Σ of $E(G)$, with $|\Sigma|$ even, the signed graph (G, Σ) is ECD.

For $X \subseteq V(G)$, we let $\delta_G(X)$ be the set of edges with exactly one end in X. We say that $\delta_G(X)$ is the *cut induced by X*. Two signatures $\Sigma_1, \Sigma_2 \subseteq E(G)$ are *equivalent* if their symmetric difference is a cut. Note that signature equivalence is (naturally) an equivalence relation. The operation of changing to an equivalent signature is called *re-signing*. The main reason for working with signed graphs is that if $\Sigma_1, \Sigma_2 \subseteq E(G)$ are equivalent signatures, then (G, Σ_1) and (G, Σ_2) have exactly the same set of even cycles, since a cycle crosses every cut an even number of times. Thus, for equivalent signatures Σ_1 and Σ_2, (G, Σ_1) is ECD if and only if (G, Σ_2) is ECD.

We will frequently use the following well-known lemma without explicit reference:

Lemma 2.2 (Folklore)**.** *Let (G, Σ) be a signed graph. For any $F \subseteq E(G)$ which does not contain a cycle, there is a signature Σ_F which is equivalent to Σ and such that $\Sigma_F \cap F = \emptyset$.*

Lemma 2.2 can be proven easily by induction on $|F|$.

We have previously noted that K_5 is not strongly ECD. On the other hand, up to re-signing, there is only one even size signature Σ of $E(K_5)$ such that (K_5, Σ) is not ECD.

Lemma 2.3. *Let Σ be a signing of K_5 where $|\Sigma|$ is even. If (K_5, Σ) is not ECD, then Σ and $E(K_5)$ are equivalent.*

Proof. By Lemma 2.2, we may assume that the edges incident with a vertex $u \in V(K_5)$ are all even. Since the complement of a 5-cycle in K_5 is another 5-cycle, every 5-cycle must be odd, else (K_5, Σ) is ECD. Every 4-cycle in K_5 is the symmetric difference of two 5-cycles, and thus is even. Every 3-cycle is the symmetric difference of a 4-cycle and a 5-cycle, and thus is odd. This means all the edges in $G - u$ are odd. Therefore, Σ and $E(K_5)$ are equivalent since their symmetric difference is $\delta(\{u\})$. $\square$

The *parity* of a vertex v in a signed graph is the parity of the number of odd non-loop edges incident with v. Note that in an Eulerian signed graph, the parity of a vertex is invariant under re-signing. Finally, if it is clear which signature Σ (up to re-signing) is under consideration, we will often abbreviate (G, Σ) by G.

3. COMPLETE BIPARTITE GRAPHS

In this section, we prove that Eulerian complete bipartite graphs are strongly ECD. In fact, we will need to prove something slightly stronger. Namely, after removing the edges of a 4-cycle from an Eulerian complete bipartite graph, the resulting graph is strongly ECD.

Let us write $K_{n,m} - C_4$ to denote a subgraph of $K_{n,m}$ obtained by deleting the edges of a fixed 4-cycle of $K_{n,m}$. We proceed via a sequence of lemmas.

Lemma 3.1. *$K_{2,n}$ is strongly ECD for all positive even integers n.*

Proof. Consider a signing Σ of $E(K_{2,n})$ with $|\Sigma|$ even. Let $\{u, v\}$ and $\{x_1, \dots, x_n\}$ be the bipartition of $K_{2,n}$. By re-signing, we may assume that all edges incident with u are even. By re-indexing vertices if necessary, we may assume that there exists an even index i such that the edges $vx_1, \dots, vx_i$ are all odd and all the other edges incident with v are even. But now,

$$\{ux_1vx_2, ux_3vx_4, ux_5vx_6, \dots, ux_{n-1}vx_n\}$$

is an even cycle decomposition of $(K_{2,n}, \Sigma)$. $\square$

Lemma 3.2. *$K_{4,4} - C_4$ is strongly ECD.*



Figure 1. Three possible signatures for $K_{4,4} - C_4$, and corresponding even cycle decompositions. Odd edges are solid and even edges are dashed.

Proof. Let $G := K_{4,4} - C_4$ with bipartition $\{a, b, c, d\}$ and $\{w, x, y, z\}$ such that $C_1 := aybz$, $C_2 := cydz$, and $C_3 := wcxd$ are edge-disjoint 4-cycles. Let Σ be a signing of $E(G)$ with $|\Sigma|$ even. If either C_1 or C_3 is even then (G, Σ) is ECD by Lemma 3.1. So assume C_1 and C_3 are both odd and therefore C_2 is even.

Note that $E(C_1)$, $E(C_2)$, and $E(C_3)$ are disjoint edge cuts. We may therefore assume by re-signing that each of C_1 and C_3 contains exactly one odd edge and that C_2 has either zero or two odd edges. Assume without loss of generality that the edges ay and cw are odd. There are three cases to consider: C_2 has no odd edges, C_2 has two odd edges forming a matching (which we can choose, by re-signing), or C_2 has two odd edges not forming a matching, which can be chosen to meet at y or c by re-signing. The even cycle decompositions for each of these cases are shown in Figure 1. □

Lemma 3.3. *$K_{n,m} - C_4$ is strongly ECD for all even integers $n, m \geq 2$.*

Proof. We proceed by induction on $n+m$. Note, that $K_{2,j} - C_4$ is $K_{2,j-2}$ (together with two isolated vertices), so by Lemma 3.1 we may assume that $n, m \geq 4$. Next, by Lemma 3.2, we may assume that $m \geq 6$. Now let $X := \{x_1, \ldots, x_n\}$ and $\{y_1, \ldots, y_m\}$ be a bipartition of $K_{n,m} - C_4$, and let Σ be a signing of its edges with $|\Sigma|$ even. By re-indexing, we may assume that the vertices of the missing C_4 are $\{x_1, x_2, y_1, y_2\}$. Since $m \geq 6$, two vertices (say, y_{m-1} and y_m) must have the same parity. Thus, the subgraphs induced by $X \cup \{y_{m-1}, y_m\}$ and $X \cup \{y_1, \ldots, y_{m-2}\}$ both contain an even number of odd edges. But now we are finished since the first subgraph is isomorphic to $K_{2,n}$ (and hence is ECD by Lemma 3.1), while the second is isomorphic to $K_{n,m-2} - C_4$ (and is ECD by induction). □

Lemma 3.4. *$K_{n,m}$ is strongly ECD for all even integers $n, m \geq 2$.*

Proof. By considering parities of vertices, it is easy to show that every even size signing Σ of such a graph must contain an even C_4. So, we remove the edges of an even C_4 and apply the previous lemma. □



FIGURE 2. A co-claw

4. ODD EXPANSION

Let G and H be simple graphs and let $v \in V(G)$. The graph obtained from the disjoint union of $G - v$ and H by joining each vertex in H to each vertex in the neighborhood of v (in G) is called *the substitution of v by H*. We may assume that this graph contains both G and H as a subgraph by identifying v with a vertex in H. In the special case that H is an odd size stable set, we refer to this operation as an *odd expansion.* (A set of vertices is called *stable* if no two vertices in the set are adjacent.)

Lemma 4.1. *If G is a simple strongly ECD graph, $v \in V(G)$, and X is an odd stable set, then the substitution of v by X is also strongly ECD.*

Proof. Let G' be the substitution of v by an odd stable set X. Let Σ be a signing of G' with $|\Sigma|$ even. Let $G_1 := G' - X$ and let $\Sigma_1 := \Sigma \cap E(G_1)$. Since $|\Sigma|$ is even and $|X|$ is odd, there exists a vertex $t \in X$ such that the parity of t is equal to the parity of $|\Sigma_1|$. Therefore, $G' - (X - t) \simeq G$ has an even number of odd edges, and hence is ECD as a signed graph. The signed graph of remaining edges is a complete bipartite graph, and is ECD by Lemma 3.4. ☐

5. ADDING APEX VERTICES

Let G and H be graphs. The *join of G and H* is the graph obtained from the disjoint union of G and H by joining each vertex in G to each vertex in H. In the special case that H has no edges, we say that the join of G and H is obtained from G by *adding apex vertices.*

An *almost even cycle decomposition* of a signed graph (G, Σ) is a partition of $E(G)$ into cycles exactly one of which is odd.

Lemma 5.1. *If G is a strongly ECD graph and Σ is a signing of G with $|\Sigma|$ odd, then (G, Σ) admits an almost even cycle decomposition.*

Proof. We flip the sign of a fixed edge and then apply the fact that G is strongly ECD. ☐

Lemma 5.2. *If G is a simple strongly ECD graph with an even number of vertices, then the join of G with an even stable set X of size at least 4 is also strongly ECD. Moreover, provided that G is not a co-claw (Figure 2), then the join of G with a stable set X of size 2 is also strongly ECD.*

Proof. Before starting the proof, we note that the join of a co-claw and a stable set of size 2 is a subdivision of K_5. Therefore, the conditions of the lemma are both necessary and sufficient.

Let G be a counterexample with $|E(G)|$ minimum, let X be an even stable set, and let G' be the join of G and X. Let Σ be a signing of G' with $|\Sigma|$ even. If

$|\Sigma \cap E(G)|$ is even as well, then $G' \setminus E(G)$ is ECD by Lemma 3.4, and we are done. Thus, $|\Sigma \cap E(G)|$ is odd. By Lemma 5.1, $(G, \Sigma \cap E(G))$ admits an almost even cycle decomposition. Note that if $|V(G)| = 4$, G has at most one cycle and removing it cannot leave us with a co-claw. Thus if the almost even cycle decomposition of $(G, \Sigma \cap E(G))$ contained an even cycle, its removal would contradict the minimality of G. We may therefore assume that $E(G) = E(C)$, where C is an odd cycle. Let $S := V(G) \setminus V(C)$.

By Lemma 2.2, we may further assume that there exists $t \in X$ such that all edges incident with t are even. Suppose the join of a pair of vertices in X and a pair of vertices in C is an odd 4-cycle D. Then $C \cup D$ can be decomposed into two even cycles. Also, $(G' \setminus E(C \cup D), \Sigma \setminus E(C \cup D))$ is ECD by Lemma 3.3.

We may hence assume that such a 4-cycle D does not exist. Therefore, for each apex vertex $x \in X$, the edges from x to C are either all even or all odd. By re-signing, we may assume they are all even. This implies that there must exist a vertex $u \in S$ of odd parity, as $|\Sigma|$ is even. We have two cases based on the size of X.

Case 1. $|X| = 2$. In this case, by hypothesis, we may assume that G is not a co-claw. Let a and b be the two vertices in X. Since u has odd parity, the edges ua and ub have different signs. We first suppose that C has length at least 4. We choose two vertices $x, y \in C$ such that one of the x-y paths in C, say P_1, is even with at most two edges. Since C has length at least 4, the other x-y path P_2 has an internal vertex z. But then $D_1 = \{xb, bz, za, ay\} \cup E(P_1)$ and $D_2 = \{xa, au, ub, by\} \cup E(P_2)$ are even cycles. By Lemma 3.4, $(G' \setminus E(D_1 \cup D_2), \Sigma \setminus E(D_1 \cup D_2))$ is ECD, and hence (G', Σ) is ECD.

We may therefore assume that C is a triangle. Since G is not a co-claw, we also have $|S| \geq 3$. Let c and d be the endpoints of an odd edge of C and let C_1 be the join of $\{c, d\}$ and X. We first suppose that there exist s_1 and s_2 in S with different parities. By symmetry we may assume that the path $P_1 := as_1b$ is odd and the path $P_2 := as_2b$ is even. Recall that we have re-signed so that all edges between X and C are even. In particular, all edges in C_1 are even. Therefore, $D_1 := \{e\} \cup \{ca, db\} \cup E(P_1)$ and $D_2 := E(C \cup C_1 \cup C_2) \setminus E(D_1)$ is an even cycle decomposition of $(C \cup C_1 \cup C_2, \Sigma \cap E(C \cup C_1 \cup C_2))$. The remaining signed graph, $(G' \setminus E(C \cup C_1 \cup C_2), \Sigma \setminus E(C \cup C_1 \cup C_2))$ is ECD by Lemma 3.4.

The only remaining possibility is if all vertices in S have the same parity. Therefore, the join of any two vertices in S and X is an even 4-cycle. By continually removing these 4-cycles, we may assume $|S| = 3$. Now, by suppressing the degree-2 vertices, there are only two cases to consider by symmetry. Namely, all edges in C are odd, or exactly one edge in C is odd. Even cycle decompositions for both these possibilities are shown in Figure 3.

Case 2. $|X| \geq 4$. By applying Lemma 4.1 to the previous case, it suffices to assume $C \simeq K_3$, $S = \{u\}$ and $X = \{a, b, c, d\}$. Again recall that C is odd, and we have re-signed so that all edges between C and X are even. Let xy be an odd edge in C. Since the paths $P_1 := aub$ and $P_2 := cud$ have different parities, $C \cup P_1 \cup \{xa, yb, xc, yd\} \cup P_2$ can be partitioned into two even cycles. We are done since the signed graph of remaining edges of G' is Eulerian with no odd edge, and hence is ECD. $\square$



FIGURE 3. The upper triangle corresponds to the cycle C. Solid edges are odd and dashed edges are even. The labelled edges illustrate the union of even cycles (three on the left and two on the right) whose removal leaves an even 4-cycle.

Lemma 5.3. *Let G be a simple strongly ECD graph such that $|V(G)|$ is odd and G is not isomorphic to K_3. Then the join of G and K_2 is strongly ECD.*

Proof. Let G' be the join of G and K_2. Let G'' be the join of $G \cup \{u\}$ and a stable set of size 2, where u is a new isolated vertex added to G. Since G'' is a subdivision of G', it follows that G' is strongly ECD if and only if G'' is strongly ECD. Thus, provided $G \not\cong K_3$, by Lemma 5.2, G' is strongly ECD. $\square$

6. SUBSTITUTING A VERTEX

Lemma 6.1. *If G is a simple strongly ECD graph, $v \in V(G)$, and H is a simple strongly ECD graph with an odd number of vertices, then the substitution of v by H is strongly ECD provided that H is not a triangle or v has degree at least 4.*

Proof. Let G' be the substitution of v by H. Let Σ be a signing of G' with $|\Sigma|$ even. If $|\Sigma \cap E(H)|$ is even, then we are done by Lemma 4.1 since $G' \setminus E(H)$ is an odd expansion of G. Thus, $|\Sigma \cap E(H)|$ is odd. By taking an almost even cycle decomposition of H, we may further assume that the signed graph $(H, \Sigma \cap E(H))$ is the disjoint union of an odd cycle C and a stable set S. Since $|V(H)|$ is odd, there exists a vertex $u \in V(H)$ such that one of the following holds.

(1) $u \in S$ and $G' \setminus V(H - u)$ has an even number of odd edges.
(2) $u \in S$ and $G' \setminus V(H - u)$ has an odd number of odd edges.
(3) $S = \emptyset$ and $G' \setminus V(H - u)$ has an odd number of odd edges.

For simplicity, we refer to $G' \setminus V(H - u)$ as G by identifying u and v. Define $G'' := G' \setminus E(G)$. Since G is strongly ECD, we can remove even cycles from $E(G)$ until just a single cycle $C' := \{ua, bu\} \cup P$ remains, where P is a path in $G - u$ joining two neighbors a, b of u in G. Thus we may assume that $E(G) = E(C')$. By re-signing, we may further assume that all edges incident with u are even. Now we show that in all three cases (G', Σ) is ECD.

Case (1). Note that G'', after removing its isolated vertices, is the join of $H - u$ and an even sized stable set. If the degree of v in G is at least 4, then we are done since $(G'', \Sigma \setminus E(G))$ is ECD by Lemma 5.2. Thus we may assume that the degree of v is 2. In this case, after removing isolated vertices in G and suppressing degree-2 vertices in P, we deduce that G' is the join of H with K_2. Observe that $H \not\simeq K_3$ since it contains the isolated vertex u. By Lemma 5.3, (G', Σ) is ECD.

Case (2). Here C' is an odd cycle. Since C has at least three vertices, there are vertices z and w in C such that $C'' := azbw$ is an even 4-cycle. Thus, $C \cup C' \cup C''$ can be decomposed into two even cycles. We are done with this case since by Lemma 3.3, $(G' \setminus E(C \cup C' \cup C''), \Sigma \setminus E(C \cup C' \cup C''))$ is ECD.

Case (3). Here $u \in V(C)$, P is odd, and S is empty. For distinct $v, w \in V(C) \setminus \{u\}$ we let $K(v,w)$ be the subgraph induced by the edges in $E(C) \cup E(C') \cup \{va, vb, wa, wb\}$. Note that $K(v,w)$ is a subdivision of K_5. For each $v \in V(C)$ we define $p(v)$ to be the parity of $|\Sigma \cap \{va, vb\}|$. We first assume that there exist distinct $v, w \in V(C) \setminus \{u\}$ such that both $p(v)$ and $p(u)$ are odd. Since $K(v,w)$ contains a (subdivided) even triangle, it is ECD by Lemma 2.3. Since u is an isolated vertex in $G' \setminus E(K(v,w))$, we have that $(G' \setminus E(K(v,w)), \Sigma \setminus E(K(v,w)))$ is ECD by Lemma 3.3.

We may hence assume that there is at most one vertex $w \in V(G) \setminus \{u\}$ such that $p(w)$ is odd. Since C is not a triangle, it has length at least 5. Therefore, there are distinct vertices $x, y, z \in V(C) \setminus \{u\}$ such that $p(x)$, $p(y)$, $p(z)$ are even. It is easy to verify that one of $K(x,y), K(x,z)$ of $K(y,z)$ (say $K(x,y)$) must contain an even (subdivided) triangle, and is hence ECD by Lemma 2.3. Again, $(G' \setminus E(K(x,y)), \Sigma \setminus E(K(x,y)))$ is ECD by Lemma 3.3. □

7. Substituting twin vertices

Two distinct vertices u and v of a simple graph G are *twins* if u and v are non-adjacent and have the same neighborhood. If u and v are twins of G and H is another graph, then *the twin substitution of* $\{u, v\}$ *by* H is the graph obtained from the disjoint union of $G - \{u, v\}$ and H by connecting each vertex in H to each neighbor of u in G.

Lemma 7.1. *Let G be a simple strongly ECD graph and let u and v be twin vertices in G. If H is a simple strongly ECD with an even number of vertices, then the twin substitution of $\{u, v\}$ by H is also strongly ECD, provided that H is not a co-claw or v has degree at least 4.*

Proof. Let G' be the twin substitution of $\{u, v\}$ by H. Let Σ be a signing of G' with $|\Sigma|$ even. If H has an even number of odd edges, then we are done by Lemma 4.1. Otherwise, by taking an almost even cycle decomposition, we may assume that H is an odd cycle plus some isolated vertices. Suppose H has an isolated vertex, say w. Then the twin substitution of $\{u, v\}$ by H is the same as substitution of u by $H - w$. Since H is not a co-claw, by Lemma 6.1, we may assume that H is an odd cycle of even length. Hence, there exist two vertices in H, say x and y, such that $(G' - H) \cup \delta(\{x, y\}, N(u)) \simeq G$ has an odd number of odd edges. (Here, $N(u)$ denotes the set of neighbors of u in G.) Fix $z \in N(u)$, and let G_0 be the signed graph obtained by flipping the sign of the edge xz in $(G' - H) \cup \delta(\{x, y\}, N(u))$. Let $\mathcal{C}_0$ be an even cycle decomposition of G_0, and let C_1 and C_2 be cycles in $\mathcal{C}_0$ containing xz and yz respectively (possibly $C_1 = C_2$). Note that in $(G' - H) \cup \delta(\{x, y\}, N(u))$, C_1 is odd and x and y are non-adjacent. Therefore, $C_1 \cup C_2$ can be partitioned into two edge disjoint x-y paths of different parities. It follows that $H \cup C_1 \cup C_2$ can be partitioned into two even cycles, and thus $H \cup E(\mathcal{C}_0)$ is ECD. The signed graph of remaining edges is ECD by Lemma 3.4. □

8. Joins and Apex Joins

Lemma 8.1. *Let G be a disjoint union of strongly ECD graphs. The join of G and a stable set of size two is strongly ECD, provided that $|V(G)|$ is even and G is not a co-claw.*

Proof. Let G' be the join of G and the stable set $\{u, v\}$. Suppose G consists of n odd connected components and m even connected components. Thus G' can be obtained from $K_{2,n+2m}$ by n substitutions and m twin substitutions. We proceed by induction on N, the number of components in G isomorphic to K_3. If $N = 0$, then G' is strongly ECD by Lemmas 6.1 and 7.1. Suppose $N \geq 1$. Let Σ be a signing of $E(G')$ with $|\Sigma|$ even. Let $H_1, \dots, H_{n+m}$ be the components of G. By considering even (or almost even) cycle decompositions of each H_i, we may assume that each component is an isolated vertex or an odd cycle. At least one component, say C_1, is odd and isomorphic to K_3, else $N = 0$. By Lemma 5.2, we may assume that another component, say C_2, is also an odd cycle. By a simple parity argurment, for each $i = 1, 2$, there are vertices x_i, y_i in C_i such that $C_i' := ux_ivy_i$ is an even 4-cycle. Then $C_1 \cup C_2 \cup C_1' \cup C_2'$ can be decomposed into two even cycles, and hence its removal from G' yields the join of $\{u, v\}$ and a graph with smaller N. ☐

Lemma 8.2. *The join of two co-claws is strongly ECD.*

Proof. Let G be the join of two co-claws. Let C_1 and C_2 be the components in the co-claws isomorphic to K_3. Let Σ be a signing of G with $|\Sigma|$ even. We may assume that C_1 and C_2 are both odd, else we can apply Lemma 5.2. Let x_1 and y_1 be two distinct vertices in C_1. By an easy parity argument, there are vertices x_2 and y_2 in C_2 such that $C := x_1x_2y_1y_2$ is an even 4-cycle. Then $C \cup C_1 \cup C_2$ can be decomposed into even cycles. The remaining edges of G can be decomposed into even cycles by Lemma 3.3. ☐

Corollary 8.3. *If G_1 and G_2 are both disjoint unions of simple strongly ECD graphs with $|V(G_1)|$ and $|V(G_2)|$ both even, then the join of G_1 and G_2 is also strongly ECD, provided that it is not the join of a co-claw and a stable set of size two.*

Proof. If either G_1 or G_2 is a stable set of size 2, then we are done by Lemma 8.1. Thus, $|V(G_1)|, |V(G_2)| \geq 4$. Suppose that G_1 consists of o_1 odd connected components and e_1 even connected components, and G_2 consists of o_2 odd connected components and e_2 even connected components. Since $|V(G_1)|$ and $|V(G_2)|$ are both even, note that o_1 and o_2 are also both even. Let G be the join of G_1 and G_2. Then G can be obtained by starting from $K_{o_1+2e_1,o_2+2e_2}$ and performing $o_1 + o_2$ substitutions and $e_1 + e_2$ twin substitutions. Now, if $o_1 + 2e_1 \geq 4$ or $o_2 + 2e_2 \geq 4$, then we are done by Lemmas 8.1, 6.1 and 7.1. Thus, $o_1 + 2e_1 = 2 = o_2 + 2e_2$. By Lemma 8.2, we may assume that G_1 is not a co-claw. Let G_1' be the join of G_1 with a stable set of size two. By Lemma 8.1, G_1' is strongly ECD. But now the join of G_1 and G_2 can be obtained from G_1' by two substitutions or a twin substitution, and is hence strongly ECD by Lemma 6.1 or Lemma 7.1. ☐

Lemma 8.4. *Let G and H be simple strongly ECD graphs with an odd number of vertices. If $|V(G)| \geq 5$, then the graph obtained by adding an apex vertex to the join of G and H is also strongly ECD.*

Proof. Let u and v be distinct vertices of K_3. Since $|V(G)| \geq 5$, substituting G for u in K_3 yields a strongly ECD graph G' by Lemma 6.1. Since v now has degree at least six in G', substituting H for v in G' yields another strongly ECD graph G''. But G'' is precisely the join of G and H together with an apex vertex. □

9. Our Main Theorem

In this section we derive Theorem 1.3 as a corollary of our composition operations.

Lemma 9.1. *If n is odd, then K_n is strongly ECD unless $n = 5$.*

Proof. Clearly K_1 and K_3 are both strongly ECD, so by Lemma 5.3 it suffices to show that K_7 is strongly ECD. To see this, start with K_3. By substituting a vertex of K_3 with a stable set of size 3, it follows that K_5 minus the edges of a triangle is strongly ECD. Again by Lemma 5.3, we have that K_7 minus the edges of a triangle is strongly ECD. Now consider a signing Σ of $E(K_7)$ with $|\Sigma|$ even. It suffices to show that (K_7, Σ) must contain an even triangle. To see this, re-sign so that each edge of a spanning star T is even. Since $|E(T)| = 6$, $|E(K_7)| = 21$, and $|\Sigma|$ is even, there must exist another even edge $e \notin E(T)$. Thus, $T \cup e$ contains an even triangle, as required. □

We finish with the promised proof of Theorem 1.3.

Theorem 1.3. *Every Eulerian complete r-partite graph is strongly ECD, except for K_5.*

Proof. Let $K = K_{n_1,\ldots,n_r}$ be a complete Eulerian r-partite graph. First observe that if each n_i is even, then K is strongly ECD since we may build K by starting from K_{n_1,n_2} and repeatedly applying Lemma 5.2. The remaining case is if some n_j is odd. Since G is Eulerian, it must be the case that all n_i are odd and r is odd. By Lemma 9.1, we have that K_n is strongly ECD for all odd $n \neq 5$. Thus, by substituting stable sets for vertices of K_r, we are done unless $r = 5$. For $r = 5$, it suffices to show that $K_{3,1,1,1,1}$ is strongly ECD. To see this we apply Lemma 8.4 with $G = K_{3,1,1}$ and $H = K_1$. □

References

[1] H. Fleischner. (Some of) the many uses of Eulerian graphs in graph theory (plus some applications). *Discrete Math.*, 230(1-3):23–43, 2001. Paul Catlin memorial collection (Kalamazoo, MI, 1996).

[2] T. Huynh, S. Oum, and M. Verdian-Rizi. Even cycle decompositions of graphs with no odd-K_4-minor. In preparation, 2012.

[3] B. Jackson. On circuit covers, circuit decompositions and Euler tours of graphs. In *Surveys in combinatorics, 1993 (Keele)*, volume 187 of *London Math. Soc. Lecture Note Ser.*, pages 191–210. Cambridge Univ. Press, Cambridge, 1993.

[4] K. Markström. Even cycle decompositions of 4-regular graphs and line graphs. *Discrete Math.*, 312(17):2676–2681, 2012.

[5] P. D. Seymour. Even circuits in planar graphs. *J. Combin. Theory Ser. B*, 31(3):327–338, 1981.

[6] C. Q. Zhang. On even circuit decompositions of Eulerian graphs. *J. Graph Theory*, 18(1):51–57, 1994.

(Tony Huynh, Sang-il Oum, Maryam Verdian-Rizi) Department of Mathematical Sciences, KAIST, 291 Daehak-ro Yuseong-gu Daejeon, 305-701 South Korea

(Andrew D. King) Department of Mathematics, Simon Fraser University, 8888 University Drive, Burnaby, BC, V5A 1S6, Canada
E-mail address: `tony.bourbaki@gmail.com`
E-mail address: `andrew.d.king@gmail.com`
E-mail address: `sangil@kaist.edu`
E-mail address: `mverdian@gmail.com`